

MARKOVIAN LOG-SUPERMODULARITY, AND ITS APPLICATIONS IN PHYLOGENETICS

MIKE STEEL AND BEATA FALLER

ABSTRACT. We establish a log-supermodularity property for probability distributions on binary patterns observed at the tips of a tree that are generated under any 2-state Markov process. We illustrate the applicability of this result in phylogenetics by deriving an inequality relevant to estimating expected future phylogenetic diversity under a model of species extinction. In a further application of the log-supermodularity property, we derive a purely combinatorial inequality for the parsimony score of a binary character. The proofs of our results exploit two classical theorems in the combinatorics of finite sets.

1. INTRODUCTION

Finite-state Markov processes on trees are widely used in evolutionary biology to model the way in which discrete characteristics of present-day species have evolved from the state present in some common ancestor [6, 12]. In this paper, we investigate a generic inequality that applies to 2-state Markov processes on trees, and provide two applications.

The first application, which was the motivation for our study, is to the theory of biodiversity conservation. We consider the expected loss of ‘phylogenetic diversity’ under a model in which extinction risk is associated with an underlying state that evolves on the tree. We are interested in comparing this expected loss to simpler models in which extinction events are treated independently; we find that that when extinction events reflect phylogenetic history, then the expected loss of phylogenetic diversity is always greater than or equal to that predicted by an independent extinction scenario. Essentially, this is because the probability that an entire ‘clade’ (the set of present-day species descended from a vertex in the tree) becomes extinct is higher when the evolution of the influencing state is taken into consideration than when we treat extinctions as independent events.

In a second application, we derive a new, purely combinatorial result concerning the ‘parsimony score’ of a binary character on a tree. We also briefly discuss how the generic inequality for 2-state Markov processes relates to recent work on phylogenetic invariants and inequalities for particular submodels.

1991 *Mathematics Subject Classification.* 05C05; 92D15.

Key words and phrases. tree, Markov process, FKG inequality, phylogenetic diversity.

We thank the New Zealand Marsden Fund for supporting this work.

2. MARKOV PROCESSES ON TREES

Let $T = (V_T, E_T)$ be a tree with leaf set X . Consider a Markov random field on T with state space $\{0, 1\}$, and for each vertex v of T , let $\xi(v)$ be the random state (0 or 1) that v is assigned. This process is usually described as follows. We have a root vertex ρ for which we specify a probability, say π_i , that $\xi(\rho) = i$, for $i \in \{0, 1\}$. Direct all the edges of T away from ρ and for any arc (r, s) of the resulting directed tree $T = (V_T, A_T)$, let $P^{(r,s)}$ denote the 2×2 transition matrix for which the ij -entry (for $i, j \in \{0, 1\}$) is the conditional probability that $\xi(s) = j$ given that $\xi(r) = i$. Specifying $\pi = [\pi_0, \pi_1]$ together with the transition matrices $P^{(r,s)}$ for all the arcs (r, s) of T uniquely defines the Markov random field on T (see, for example, [3, 12, 14]); an explicit formula appears below (Eqn. 1). We will assume throughout that π is strictly positive and that the following condition holds on each of the transition matrices:

$$\det P^{(r,s)} \geq 0.$$

Notice that this determinant condition automatically holds if one views the transition matrix for an arc as describing the net effect of a continuous-time Markov process operating for some duration for that arc. Note however that we are not assuming that any such process is the same between the arcs of T (i.e. the model is not necessarily stationary).

For $U \subseteq V_T$ let $P(U)$ denote the probability that U is precisely the set of vertices of T in state 0; that is:

$$P(U) = \mathbb{P}(\{v \in V_T : \xi(v) = 0\} = U).$$

To express $P(U)$ in terms of the transition matrices and π , let:

$$\delta(U, v) = \begin{cases} 0, & \text{if } v \in U; \\ 1, & \text{if } v \in V_T - U. \end{cases}$$

Then, the Markov property gives:

$$(1) \quad P(U) = \pi_{\delta(U, \rho)} \cdot \prod_{(r,s) \in A_T} P_{\delta(U, r)\delta(U, s)}^{(r,s)}.$$

For any subset W of X (the leaf set of T), let p_W denote the probability that W is precisely the set of leaves of T that are in state 0. This marginal probability is given by:

$$(2) \quad p_W = \sum_{U \in \mathcal{A}_W} P(U)$$

where:

$$(3) \quad \mathcal{A}_W := \{U \subseteq V_T : U \cap X = W\}.$$

An example to illustrate this concept is provided in Fig. 1.

A number of authors have noticed that certain inequalities hold for quadratic functions of the p_W values. For example, for any $x, y \in X$ with $x \neq y$, it is well known that:

$$p_{\{x\}} \cdot p_{\{y\}} \leq p_{\{x,y\}} \cdot p_{\emptyset}.$$




FIGURE 1. In this example, if $W = \{a, c\}$, then $p_W = \pi_0 P_{00}^{(\rho,a)} P_{01}^{(\rho,b)} P_{00}^{(\rho,s)} P_{00}^{(s,c)} P_{01}^{(s,d)} + \pi_0 P_{00}^{(\rho,a)} P_{01}^{(\rho,b)} P_{01}^{(\rho,s)} P_{10}^{(s,c)} P_{11}^{(s,d)} + \pi_1 P_{10}^{(\rho,a)} P_{11}^{(\rho,b)} P_{10}^{(\rho,s)} P_{00}^{(s,c)} P_{01}^{(s,d)} + \pi_1 P_{10}^{(\rho,a)} P_{11}^{(\rho,b)} P_{11}^{(\rho,s)} P_{10}^{(s,c)} P_{11}^{(s,d)}$

Moreover, in [11] the following inequality was described: for subsets $\{x, y\}$ and $\{x, z\}$ of X where x, y, z are distinct, we have

$$P_{\{x,y\}} P_{\{x,z\}} \leq P_{\{x,y,z\}} P_{\{x\}}.$$

The following proposition shows that these are special cases of a much more general inequality.

Proposition 2.1. *For any 2-state Markov process on a tree with leaf set X , and any two subsets Y, Z of X , we have:*

$$p_Y \cdot p_Z \leq p_{Y \cup Z} \cdot p_{Y \cap Z}.$$

Proof. Let A, B be arbitrary subsets of V_T . We first establish the following:

$$(4) \quad P(A) \cdot P(B) \leq P(A \cup B) \cdot P(A \cap B).$$

Applying Eqn. (1) to $U \in \{A, B, A \cup B, A \cap B\}$, the product $P(A) \cdot P(B)$ and the product $P(A \cup B) \cdot P(A \cap B)$ can each be written as a product of two entries of π multiplied by a product over the arcs (r, s) of T of two entries of $P^{(r,s)}$. Moreover, regardless of where r and s lie in relation to the sets A, B , the product of the two π terms agree in $P(A) \cdot P(B)$ and $P(A \cup B) \cdot P(A \cap B)$ (i.e., we have $\pi_{\delta(A,\rho)} \pi_{\delta(B,\rho)} = \pi_{\delta(A \cup B, \rho)} \pi_{\delta(A \cap B, \rho)}$), while the product of the two $P^{(r,s)}$ terms agree in $P(A) \cdot P(B)$ and $P(A \cup B) \cdot P(A \cap B)$, except for the cases in which either (i) $r \in A - B$ and $s \in B - A$, or (ii) $r \in B - A$ and $s \in A - B$. However, in both case (i) and (ii), the product $P_{01}^{(r,s)} P_{10}^{(r,s)}$ appears in the term for $P(A) \cdot P(B)$ while $P_{00}^{(r,s)} P_{11}^{(r,s)}$ appears in the term for $P(A \cup B) \cdot P(A \cap B)$, and the former term is less or equal to the second since

$$P_{00}^{(r,s)} P_{11}^{(r,s)} - P_{01}^{(r,s)} P_{10}^{(r,s)} = \det P^{(r,s)}$$

and $\det P^{(r,s)} \geq 0$ by assumption. Consequently, all the terms in $P(A) \cdot P(B)$ are either less than or equal to (in cases (i) and (ii)), or equal to (in all remaining cases) the corresponding terms in $P(A \cup B) \cdot P(A \cap B)$. This establishes (4).

We now invoke a classical result of Ahlswede and Daykin (1978) [1], sometimes called the ‘four functions theorem’. A particular form of this theorem that suffices for our purposes is the following (we follow [2]). Suppose we have a finite set S and a function α that assigns a non-negative real number to each subset of S . Suppose that α satisfies the property that for all subsets A, B of S

$$\alpha(A) \alpha(B) \leq \alpha(A \cup B) \alpha(A \cap B).$$

For a collection $\mathcal{C}$ of subsets of S , let $\alpha(\mathcal{C}) := \sum_{C \in \mathcal{C}} \alpha(C)$. Then for any two collection of subsets of S , $\mathcal{A}$ and $\mathcal{B}$, say, we have:

$$(5) \quad \alpha(\mathcal{A})\alpha(\mathcal{B}) \leq \alpha(\mathcal{A} \vee \mathcal{B})\alpha(\mathcal{A} \wedge \mathcal{B}),$$

where

$$\begin{aligned} \mathcal{A} \vee \mathcal{B} &:= \{E \subseteq S : E = A \cup B : A \in \mathcal{A}, B \in \mathcal{B}\}, \text{ and} \\ \mathcal{A} \wedge \mathcal{B} &:= \{E \subseteq S : E = A \cap B : A \in \mathcal{A}, B \in \mathcal{B}\}. \end{aligned}$$

We will apply this to our problem by taking $S = V_T$, $\alpha(U) = P(U)$ and noting that α satisfies the required hypothesis by (4). Recall the definition of $\mathcal{A}_W$ in (3) and note that:

$$\mathcal{A}_Y \vee \mathcal{A}_Z = \mathcal{A}_{Y \cup Z}, \text{ and } \mathcal{A}_Y \wedge \mathcal{A}_Z = \mathcal{A}_{Y \cap Z}.$$

Thus taking $\mathcal{A} = \mathcal{A}_Y$ and $\mathcal{B} = \mathcal{A}_Z$ in (5) we deduce that:

$$\alpha(\mathcal{A}_Y)\alpha(\mathcal{A}_Z) \leq \alpha(\mathcal{A}_{Y \cup Z})\alpha(\mathcal{A}_{Y \cap Z}).$$

The proposition now follows by observing that $p_W = \alpha(\mathcal{A}_W)$ for all subsets W of X , in particular the subsets $Y, Z, Y \cup Z$ and $Y \cap Z$.

□

3. APPLICATIONS IN PHYLOGENETICS

We now describe some applications of Proposition 2.1.

3.1. Expected future phylogenetic diversity. We first show how Proposition 2.1, together with another inequality, provides a general inequality concerning the loss of expected future biodiversity under species extinction models.

Suppose that T is a rooted tree with leaf set X , and with each arc $e = (u, v)$ of T there is an associated length λ_e . Given a subset Y of X , the *phylogenetic diversity* (PD) of Y , denoted φ_Y , is the sum of the lengths of the edges of the minimal subtree of T connecting the root and the leaves in Y . Under various possible interpretations of the λ values, PD has been widely used as a measure for quantifying present and expected future biodiversity [4, 5, 10].

For each species $x \in X$ let E_x denote the event that species x is extinct at some future time t . Then the expected phylogenetic diversity of the set of species that are extant at time t , referred to as *expected future PD* and denoted $\mathbb{E}[\varphi]$, is given by:

$$(6) \quad \mathbb{E}[\varphi] = \sum_{e=(u,v) \in A_T} \lambda_e \cdot (1 - \mathbb{P}(\bigcap_{x \in C_v} E_x)) = \varphi_X - \sum_{e=(u,v) \in A_T} \lambda_e \cdot \mathbb{P}(\bigcap_{x \in C_v} E_x),$$

where C_v denotes the subset of X that is separated from the root by v . A simple model, referred to as the *generalized field of bullets model* (g-FOB) in [5] (generalizing an earlier model from [10]), assumes that the events E_x are independent. Then, if we let $p_x = \mathbb{P}(E_x)$, the value of $\mathbb{P}(\bigcap_{x \in C_v} E_x)$ in (6) (the probability of the extinction of all the species descended from v) is given by:

$$(7) \quad \mathbb{P}(\bigcap_{x \in C_v} E_x) = \prod_{x \in C_v} p_x.$$

An example to illustrate this concept is provided in Fig. 2.




FIGURE 2. If the species indicated by * become extinct, then the remaining phylogenetic diversity is the sum of the lengths of the bold edges indicated.

The assumption that the events E_x are independent is likely to be unrealistic in most settings (see, for example, [8, 13]). For example, species that are ‘close together’ in T are more likely to share attributes that may put them at risk in a hostile future environment.

To take a simple but topical scenario, consider extinction risk due to climate change. Suppose that the extinction risk of each species in X is partially influenced by some associated binary state (0 or 1) where state 0 confers an elevated risk of extinction under climate change. We suppose that these states are not known in advance for the species in X , and that this state has evolved under some Markovian model on t . Once the states are determined at the leaves, then extinction proceeds according to the g-FOB model, where species x is extinct at time t with probability p_x^i if it is in state $i \in \{0, 1\}$. We call this a *state-based field of bullets* model (s-FOB). Note that this includes the g-FOB model as a special case where $p_x^0 = p_x^1$ for all x . Moreover, once we condition on the state for each leaf, an s-FOB model is just a g-FOB model with modified extinction probabilities, but we are assuming that these states are unknown (in line with the uncertainty over what features may be helpful for an organism in a future climate).

With any s-FOB model we also have an associated g-FOB model in which the extinction probability of each species x is the same as in the s-FOB model. That is, in the g-FOB model we set:

$$(8) \quad p_x = p_x^0 \mathbb{P}(\xi(x) = 0) + p_x^1 \mathbb{P}(\xi(x) = 1),$$

where ξ describes the Markov process for the binary character. A natural question arises: how does the future expected PD of an s-FOB model compare with that of its associated g-FOB model? The following result provides a general inequality.

Theorem 3.1. *Consider a fixed tree with branch lengths and leaf set X . Consider an s-FOB model, in which state 1 is advantageous for each species, i.e., $p_x^1 \leq p_x^0$ for all $x \in X$. Then the expected future PD of this model is less or equal to the expected future PD of the associated g-FOB model.*

Proof. In view of (6) and (7), it suffices to show that:

$$(9) \quad \prod_{x \in C_v} p_x \leq \mathbb{P}\left(\bigcap_{x \in C_v} E_x\right),$$

where p_x is defined by Eqn. (8).

For each subset W of C_v let p_W denote the probability that the set of elements of C_v in state 0 is precisely W . Then:

$$\mathbb{P}\left(\bigcap_{x \in C_v} E_x\right) = \sum_{W \subseteq C_v} p_W \prod_{x \in W} p_x^0 \prod_{x \in C_v - W} p_x^1.$$

Thus, if we let:

$$f_x(W) = \begin{cases} p_x^0, & \text{if } x \in W; \\ p_x^1, & \text{if } x \in C_v - W. \end{cases}$$

then:

$$\mathbb{P}\left(\bigcap_{x \in C_v} E_x\right) = \sum_{W \subseteq C_v} p_W \prod_{x \in C_v} f_x(W).$$

Moreover:

$$p_x = p_x^0 \mathbb{P}(\xi(x) = 0) + p_x^1 \mathbb{P}(\xi(x) = 1) = \sum_{W \subseteq C_v} p_W f_x(W),$$

where the second equality arises by considering in the summation those W containing x and those not containing x . Consequently, (9) is equivalent to the requirement that:

$$(10) \quad \prod_{x \in C_v} \left(\sum_{W \subseteq C_v} p_W f_x(W) \right) \leq \sum_{W \subseteq C_v} p_W \prod_{x \in C_v} f_x(W).$$

The proof of (10) involves combining Proposition 2.1 with the FKG inequality of Fortuin, Kasteleyn and Ginibre (1971) [7], a particular (and multivariate) form of which we now recall.

Given a finite set S , suppose that $f_1, f_2, \dots, f_n$ are functions from the power set of S into the non-negative real numbers, and that these satisfy the condition:

$$(11) \quad A \subseteq B \Rightarrow f_i(A) \leq f_i(B).$$

Furthermore, suppose that μ is a probability measure on the subsets of S which satisfies the log-supermodularity condition:

$$(12) \quad \mu(A)\mu(B) \leq \mu(A \cup B)\mu(A \cap B).$$

Then:

$$(13) \quad \prod_{i=1}^n \left(\sum_A \mu(A) f_i(A) \right) \leq \sum_A \mu(A) \prod_{i=1}^n f_i(A)$$

where the summations are over all subsets of S .

We apply this form of the FKG inequality by taking $S = \{1, \dots, n\} = C_v$, $\mu(W) = p_W$, and f_x as defined above. Then f_x satisfies (11) by the hypothesis that $p_x^1 \leq p_x^0$ for all x , while μ satisfies (12) by Proposition 2.1. Then inequality (13) provides the required inequality (10). This completes the proof.

□

3.2. Combinatorics of parsimony. We now describe a further application of Proposition 2.1 by deriving a purely combinatorial result concerning a measure (‘parsimony score’) that underlies certain approaches for inferring evolutionary history (see, for example, [6]).

Given a function $f : X \rightarrow \{0, 1\}$, recall that the *parsimony score* of f on T , denoted $l(f, T)$, is the minimum number of edges that have different states assigned to their endpoints, across all extensions $F : V_T \rightarrow \{0, 1\}$ of f . For example, for the tree T in Fig. 1, and the function f defined by $f(a) = f(c) = 0, f(b) = f(d) = 1$, we have $l(f, T) = 2$; for this example, there are two minimal extensions F of f corresponding to $F(\rho) = F(s) = i$ for $i \in \{0, 1\}$ (for further details concerning the mathematical properties of parsimony score, see [12]). For $W \subseteq X$, let f_W denote the function that assigns state 0 to the elements of W , and assigns state 1 to the elements of $X - W$. The following result states that the parsimony score function for a given tree is submodular.

Theorem 3.2. *For any tree T with leaf set X and subsets Y, Z , of X we have:*

$$l(f_Y, T) + l(f_Z, T) \geq l(f_{Y \cup Z}, T) + l(f_{Y \cap Z}, T).$$

Proof. Consider the 2-state Markov random field on T with $\pi_0 = \pi_1 = 0.5$, and set each transition matrix $P^{(r,s)}$ to be the symmetric 2×2 matrix with off-diagonal entry $\epsilon > 0$. Then, for any $W \subseteq X$ a straightforward calculation shows that:

$$(14) \quad p_W = C_W \epsilon^{l(f_W, T)} (1 + o(\epsilon)),$$

for a constant C_W that depends only on W and T and not ϵ (specifically, C_W is the number of minimal extensions of f_W to the vertices of T multiplied by $\frac{1}{2}$). Now Proposition 2.1, expressed using logarithms, states that:

$$(15) \quad -\log(p_Y) - \log(p_Z) \geq -\log(p_{Y \cup Z}) - \log(p_{Y \cap Z}).$$

Applying (14) (and noting that $\log(1 + o(\epsilon)) = o(\epsilon)$), the left-hand side of (15) is:

$$(l(f_Y, T) + l(f_Z, T)) \log\left(\frac{1}{\epsilon}\right) - \log(C_Y C_Z) + o(\epsilon)$$

while the right-hand side of (15) is:

$$(l(f_{Y \cup Z}, T) + l(f_{Y \cap Z}, T)) \log\left(\frac{1}{\epsilon}\right) - \log(C_{Y \cup Z} C_{Y \cap Z}) + o(\epsilon).$$

Theorem 3.2 now follows by letting ϵ tend to zero. □

4. CONCLUDING REMARKS

- (i) It is possible to establish Theorem 3.2 using a purely combinatorial proof, by first invoking Menger’s theorem from graph theory to handle the case where Y and Z are disjoint, and then using a complementation argument for the case $Y \cup Z = X$. The remaining case where $X - (Y \cup Z)$ is non-empty can then be established by a somewhat detailed argument that uses induction on $|X|$.

- (ii) Proposition 2.1 provides a collection of polynomial inequalities on the p_W values, which have recently been studied for a particular class of Markov 2-state models in [9]. These polynomial inequalities complement the much-studied ‘phylogenetic invariants’ (polynomial identities in the p_W values), which hold under various restrictions on the Markov model. Combining these phylogenetic invariants with the polynomial inequalities provides a way of characterizing when a probability distribution arises on a tree under a Markov process (either with or without restrictions). For the 2-state Markov process on a tree with 3 leaves, this was solved in [11].
- (iii) It may be of interest to derive an extension of Proposition 2.1 that applies when the state space has a size greater than 2.

REFERENCES

- [1] R. Ahlswede, and D.E. Daykin, An inequality for the weights of two families of sets, their unions and intersections, *Z. Wahrsch. V. Geb.* **43** 183-185 (1978).
- [2] I. Anderson, *Combinatorics of finite sets*, Dover Publications, New York, (1987).
- [3] J.T. Chang, Full reconstruction of Markov models on evolutionary trees: identifiability and consistency, *Math. Biosci.* **137** 51-73 (1996).
- [4] D.P. Faith, Conservation evaluation and phylogenetic diversity, *Biol. Conserv.* **61** 1-10 (1992).
- [5] B. Faller, F. Pardi and M. Steel, Distribution of phylogenetic diversity under random extinction, *J. Theor. Biol.* **251** 286-296 (2008).
- [6] J. Felsenstein, *Inferring phylogenies*, Sinauer Press, (2003).
- [7] C.M. Fortuin, P.W. Kasteleyn and J. Ginibre, Correlation inequalities on some partially ordered sets, *Commun. math. Phys.* **22** 89-103 (1971).
- [8] S.B. Heard, and A.O. Mooers, Phylogenetically patterned speciation rates and extinction risks change the loss of evolutionary history during extinctions, *Proc. R. Soc. Lond. B.* **267** 613-620 (2000).
- [9] F.A. Matsen, Fourier transform inequalities for phylogenetic trees. arXiv:0711.3492 (q-bio.PE) (2008).
- [10] S. Nee and R.M. May, Extinction and the loss of evolutionary history, *Science* **278**(5338) 692-694 (1997).
- [11] J. Pearl and M. Tarsi Structuring causal trees, *J. Complexity*, **2**, 60-77 (1986).
- [12] C. Semple and M. Steel, *Phylogenetics*, Oxford University Press, Oxford, (2003).
- [13] H. Simianer, Accounting for non-independence of extinction probabilities in the derivation of conservation priorities based on Weitzman’s diversity concept, *Conserv. Genet.* **9**, 171-179 (2008).
- [14] M. Steel Recovering a tree from the leaf colourations it generates under Markov model. *Appl. Math. Lett.*, **7** 19-23 (1994).

ALLAN WILSON CENTRE FOR MOLECULAR ECOLOGY AND EVOLUTION, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF CANTERBURY, CHRISTCHURCH, NEW ZEALAND

E-mail address: m.steel@math.canterbury.ac.nz